



## Original Investigation | Psychiatry

# Interventions to Reduce Mental Health Stigma in Young People A Systematic Review and Meta-Analysis

Marcelo A. Crockett, PhD; Daniel Núñez, PhD; Pablo Martínez, PhD; Francesca Borghero, MD, MSc; Susana Campos, MSc; Álvaro I. Langer, PhD; Jimena Carrasco, PhD; Vania Martínez, MD, PhD

## Abstract

**IMPORTANCE** Mental health stigma is a considerable barrier to help-seeking among young people.

**OBJECTIVE** To systematically review and meta-analyze randomized clinical trials (RCTs) of interventions aimed at reducing mental health stigma in young people.

**DATA SOURCES** Comprehensive searches were conducted in the CENTRAL, CINAHL, Embase, PubMed, and PsycINFO databases from inception to February 27, 2024. Search terms included "stigma," "mental health," "mental disorders," "adolescents," "youth," and "randomized controlled trial."

**STUDY SELECTION** Inclusion criteria encompassed RCTs involving interventions aimed at reducing mental health stigma among young people (aged 10-24 years). Studies had to report outcomes related to stigma or help-seeking behaviors. Exclusion criteria included grey literature and studies without results.

**DATA EXTRACTION AND SYNTHESIS** Data were extracted independently by 7 authors (M.A.C., D.N., F.B., S.C., Á.I.L., J.C., V.M.) using Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines. Risk of bias was assessed with the Cochrane risk-of-bias tool. Three-level multivariate meta-analyses were conducted to account for within-study correlations and to maximize data use. Standardized mean differences (SMDs) (Hedges *g*) and odds ratios (ORs) with 95% CIs were calculated. The data analysis was conducted from May 30 through July 4, 2024.

**MAIN OUTCOMES AND MEASURES** Primary outcomes included stigma-related knowledge, attitudes, behaviors, and general stigma. Help-seeking outcomes were categorized into attitudes, intentions, and behaviors. Secondary outcomes included self-efficacy and empowerment.

**RESULTS** A total of 97 studies were included in the systematic review, representing 43 852 young people (mean [IQR] age, 18.7 [15.8-21.3] years; mean [IQR] females, 59.2% [49.4%-72.0%]), and 74 studies were included in 3-level multivariate meta-analyses. Significant short-term effect sizes were found for stigma-related knowledge (SMD, 0.66; 95% CI, 0.43-0.89), attitudes (SMD, 0.38; 95% CI, 0.20-0.56), behaviors (SMD, 0.29; 95% CI, 0.13-0.45), and general stigma (SMD, 0.20; 95% CI, 0.06-0.34) and for help-seeking attitudes (SMD, 0.18; 95% CI, 0.09-0.28) and intentions (SMD, 0.14; 95% CI, 0.07-0.21). Social contact interventions had a greater influence on stigma-related behaviors than did educational approaches.

**CONCLUSIONS AND RELEVANCE** These findings suggest that interventions to reduce mental health stigma among youth are beneficial in the short term. Further high-quality RCTs with long-term follow-up are needed to better understand and enhance these interventions' outcomes.

JAMA Network Open. 2025;8(1):e2454730. doi:10.1001/jamanetworkopen.2024.54730

**Open Access.** This is an open access article distributed under the terms of the CC-BY License.

## Key Points

**Question** Are interventions aimed at reducing mental health stigma in young people associated with improved outcomes?

**Findings** In this systematic review and meta-analysis of 97 randomized clinical trials involving 43 852 young people (aged 10-24 years), interventions had a significant short-term association with improvements in stigma-related knowledge, attitudes, and behaviors and help-seeking attitudes and intentions.

**Meaning** These findings suggest that interventions to reduce mental health stigma may be effective in the short term, but further high-quality studies are needed to assess long-term outcomes.

## + Supplemental content

Author affiliations and article information are listed at the end of this article.

## Introduction

Stereotypes, prejudice, or discrimination against individuals with mental illness, known as mental health stigma,<sup>1</sup> present considerable barriers to help-seeking among young people with mental health conditions<sup>2,3</sup> (ie, mental disorders; psychosocial disabilities; other mental states associated with distress, impairment, or self-harm risk<sup>4</sup>). This stigma affects individuals and communities across cultures and is linked to several negative consequences in personal, structural, health, social, and economic domains.<sup>5</sup> Reducing mental health stigma is crucial, especially in youth, as untreated mental illness can have lasting effects.<sup>6-8</sup> Educational approaches and social contact with individuals with mental illness are commonly studied interventions to reduce stigma among young people.<sup>5,9</sup> These interventions have been shown to positively influence stigma-related attitudes and knowledge in the short term, with educational approaches being more effective in school-age populations<sup>10-13</sup> and social contact interventions being more effective in higher education students.<sup>14,15</sup> However, more high-quality studies with more extended follow-up periods are needed to understand the lasting impact of these interventions.<sup>5,9</sup> Previous meta-analyses on stigma interventions have not accounted for the correlation between mental health stigma measures within the same participants across studies.<sup>1,16,17</sup> Addressing this correlation may ensure accurate estimation of each outcome without conflation.<sup>18,19</sup> To fill this gap, we conducted a systematic review of randomized clinical trials (RCTs) on interventions to reduce mental health stigma in young people, using 3-level multivariate meta-analyses to account for these correlations and maximize data use.<sup>18,19</sup> In addition, differences in the effect size of interventions based on study characteristics were examined.

## Methods

This systematic review and meta-analysis followed the Preferred Reporting Items for Systematic Review and Meta-Analysis (PRISMA) guideline. The study protocol was preregistered with PROSPERO (CRD42020210901) and published.<sup>20</sup> Protocol amendments are documented in eAppendix 1 in Supplement 1. Human research ethics committee approval and informed consent were not required per the Agencia Nacional de Investigación y Desarrollo ethics guidelines because anonymized information from publicly available studies was analyzed.

## Eligibility Criteria

We searched studies in peer-reviewed journals indexed in the CENTRAL, CINAHL, Embase, PubMed, and PsycINFO databases from inception until February 27, 2024. Articles in English and Spanish were included to expand the search for articles from different contexts. We excluded grey literature, study protocols, and reports without results.

The study eligibility criteria followed the population, interventions, comparators, and outcomes framework. The population included young people aged 10 to 24 years with or without mental health problems. Interventions included those aimed at reducing mental health stigma, including educational approaches (ie, conferences, text reading, role play), social contact, famous films, or other types.<sup>15</sup> Comparators included active comparison (antistigma intervention), standard treatment, waiting list, and placebo or no intervention.

Outcomes were based on a sociocognitive model of stigma and a framework for help-seeking for mental health problems.<sup>1,21</sup> The primary outcomes were stigma-related knowledge, attitudes, behaviors, and general stigma (outcomes that measure knowledge, attitudes, and/or behaviors at the same time) and help-seeking attitudes, intentions, and behaviors further divided into formal (eg, health professionals) and informal (eg, friends) sources. Secondary outcomes included helping attitudes and behaviors, self-efficacy, and empowerment. Timing was categorized as short term (<3 months' follow-up), midterm (3-6 months' follow-up), and long term (>6 months' follow-up).

Study designs included RCTs and controlled clinical trials (ie, where randomization was not explicitly reported but cannot be ruled out). Systematic reviews and meta-analyses were examined for eligible primary studies.

### Search Strategy

The detailed search strategy is included in eAppendix 2 in [Supplement 1](#). The basic search strategy combined index and free-text terms for "stigma" AND ("mental health" OR "mental disorders") AND ("adolescents" OR "youth") AND "randomized controlled trial."

### Study Selection and Data Collection Process

Study selection and data extraction were conducted by 7 authors independently (M.A.C., D.N., F.B., S.C., Á.I.L., J.C., V.M.) and in duplicate until high agreement was reached. The authors then reviewed the remaining studies independently. The results were discussed among the reviewers.

Data were extracted between May 14, 2021, and April 26, 2022, and updated between February 27 and May 28, 2024. Data extracted included the following: (1) study identification (authors, year of publication, and country), (2) characteristics of the study sample and methods (age range, gender, presence and type of mental illness, criteria used to define stigma, study design, recruitment methods, and inclusion and exclusion criteria), (3) characteristics of interventions and comparators (periodicity, duration, format, content, target population, and type of stigma addressed), (4) characteristics of outcomes (type of outcome measure, measurement instruments, and timing), and (5) study results (means and standard deviations for continuous outcomes or number of events and total number of participants for dichotomous outcomes per each group).

### Risk of Bias in Individual Studies

The Cochrane risk-of-bias tool was used to assess the risk of selection; performance; detection; attrition; and notification bias, classified as high, low, or unclear.<sup>22</sup> This assessment followed the same procedures as study selection and data collection.

### Statistical Analysis

To account for the interdependence of stigma-related outcomes, we performed a 3-level multivariate meta-analysis, modeling within-study correlations to improve precision and reduce bias.<sup>18,19</sup> This approach allowed us to borrow strength across related outcomes, enhancing the robustness of our effect size estimates compared with univariable analyses.<sup>18,19</sup> For the meta-analysis, all articles that provided sufficient data for primary and/or secondary outcomes were included. If basic summary measures were not reported (eg, mean and standard deviation), we followed the Cochrane Collaboration's recommended procedures.<sup>23</sup> Meta-analyses considered the timing of outcome measurement, the type of outcomes, and the type of comparison as stratification variables.

Effect sizes were the standardized mean difference (SMD) (Hedges *g*) and odds ratio (OR) for continuous and dichotomous outcomes, respectively, with 95% CIs. Unique identifiers for each effect size and study were included as random effects to account for data structure. We approximated the variance-covariance matrix of dependent within-study effect sizes by assuming a strong correlation (0.75) within outcome categories or a moderate correlation (0.40) otherwise.<sup>16</sup> These meta-analyses estimated between- and within-study variance components with heterogeneity statistics ( $I^2$ ).<sup>16</sup> Forest plots were used to visually display the results of meta-analyses that included at least 10 studies.

Moderation analyses were performed using meta-regression techniques in meta-analyses of at least 10 studies to ensure statistical power,<sup>24</sup> with the aim of examining possible differences in the effect size of the interventions based on the characteristics of the studies. Univariable meta-regressions were used to evaluate selected study characteristics (ie, risk-of-bias assessment, type of intervention, intervention format, sample age).


Publication bias was assessed in meta-analyses of at least 10 studies through funnel plots and univariable 3-level meta-regression, with the standard error of effect sizes as the independent variable. All analyses were performed using the metafor package in R, version 4.2.0 (R Foundation).<sup>25</sup> The data analysis was conducted from May 30 through July 4, 2024.

Results

Study Selection and Characteristics

The study selection process concluded with the inclusion of 97 studies<sup>26-125</sup> for the systematic review and 74 studies<sup>51-125</sup> for the 3-level multivariate meta-analyses (**Figure 1**). Most studies were individual-level RCTs (62 studies [63.9%])<sup>26, 27, 31-33, 37-40, 42, 43, 45-48, 50-59, 61-63, 65, 66, 68, 69, 71, 73-75, 79, 82, 84, 86, 87, 89, 94-100, 102, 103, 106-108, 110, 112, 115-117, 119, 121, 122</sup> and came from developed countries, mainly the US (30 studies [30.9%]).<sup>26, 27, 33, 39, 41, 47, 48, 51, 56, 58, 59, 62, 71, 76, 79, 82-84, 95-100, 106, 112, 113, 115, 122, 123</sup> The included studies represented 43 852 young people (mean [range], 452 [32-6679] individuals; mean [IQR] age, 18.7 [15.8-21.3] years; mean [IQR] gender, 59.2% [49.4%-72.0%] female, 40.0% [27.7%-50.0%] male, and 0.8% [0.0%-0.0%] transgender or nonbinary or other gender) and frequently comprised college students (45 studies [46.4%]).<sup>27, 30, 32, 33, 37-41, 46, 47, 50, 51, 54-59, 61-63, 65, 68, 71, 76, 79, 82, 84, 86, 87, 89, 90, 94-97, 106, 107, 112, 118-122</sup> A total of 28 studies [28.9%] reported results of mixed interventions,<sup>28, 32, 40, 41, 57, 63, 65, 66, 72, 77, 78, 80, 83, 86, 88, 90, 92, 94, 101, 104-106, 118-121, 123, 125</sup> with social contact plus conferences (ie, a type of educational approach) being the most studied combination (7 studies [7.2%]).<sup>28, 57, 83, 86, 90, 101, 119</sup> For single interventions, educational conferences were the most studied type of intervention (16 studies [16.5%]).<sup>29, 30, 36, 59, 67, 74, 76, 81, 85, 87, 96, 97, 111, 113, 114, 116</sup> A total of 91 studies (93.8%) assessed stigma-related outcomes,<sup>26-38, 40-48, 50-60, 64-69, 71-108, 110-113, 115-121, 123-125</sup> 39 (40.2%) provided data for help-seeking outcomes,<sup>27, 32, 37, 39, 46, 50, 57, 58, 61-63, 66, 67, 69, 72, 75, 76, 88, 95-97, 99-104, 108, 110-116, 120, 122, 124, 125</sup> and 11 (11.3%) provided secondary outcomes.<sup>42, 51, 65, 81, 88, 94, 102, 106, 114, 116, 124</sup> Further details are presented in eTable 1 in Supplement 1.

Figure 1. Flow Diagram of the Study Selection Process



## Risk of Bias Within Studies

Studies were most frequently assessed with a high risk of bias on masking of participants and personnel (59 studies [60.8%])<sup>27-34, 38, 40, 42, 43, 45, 47, 48, 50, 54-56, 59, 60, 63, 64, 66-69, 74, 76-78, 80-90, 92-94, 96, 97, 102, 104-107, 111, 113, 114, 116, 117, 119, 125</sup> and masking of outcome assessment (43 studies [44.3%]),<sup>27-30, 34, 38-42, 45, 47, 56, 59-61, 64, 67, 74, 76-82, 84, 85, 87, 89, 90, 92-94, 97, 105-107, 111, 114, 118, 123, 125</sup> with an unclear risk of bias on allocation concealment (50 studies [51.5%]),<sup>26, 27, 29, 30, 32-36, 39, 40, 45-47, 52, 54-56, 59, 61, 62, 64, 67, 71, 73, 74, 76, 78, 79, 81, 82, 84, 85, 87, 93, 95-100, 105-107, 113, 115, 117, 118, 123, 124</sup> A low risk of bias was commonly found for selective outcome reporting (83 studies [85.6%]),<sup>27-29, 32, 33, 36-40, 43, 44, 48, 50-62, 64-66, 68, 69, 71-100, 102-108, 110-112, 114-125</sup> random sequence generation (68 studies [70.1%]),<sup>28-30, 32-35, 37-39, 41, 42, 45-47, 50, 51, 53-55, 58-65, 67-69, 71, 72, 74-79, 82, 85, 88-94, 97, 101-108, 110-114, 116, 117, 120-123</sup> and incomplete outcome data (57 studies [58.8%])<sup>31, 32, 34, 36, 37, 40, 42-46, 50-57, 62, 63, 65, 66, 68, 69, 71, 72, 76, 78, 79, 83, 84, 86-90, 92-95, 97, 101-103, 108, 110, 112, 114-117, 120-124</sup> (eTable 2 in [Supplement 1](#)).

## Syntheses of Results

### Outcomes Associated With Stigma

As shown in **Figure 2**, meta-analyses of continuous data for short-term effects of active vs control interventions revealed a significant medium effect size for knowledge (SMD, 0.66; 95% CI, 0.43-0.89; 20 studies;  $I^2 = 73.3\%$ )<sup>64-83</sup> and significant small effect sizes for attitudes (SMD, 0.38; 95% CI, 0.20-0.56; 27 studies;  $I^2 = 74.0\%$ ),<sup>71-97</sup> behaviors (SMD, 0.29; 95% CI, 0.13-0.45; 22 studies;  $I^2 = 75.3\%$ ),<sup>51,52,65,66,70-73,78-82,84-90,112,124</sup> and general stigma (SMD, 0.20; 95% CI, 0.06-0.34; 13 studies;  $I^2 = 56.0\%$ ).<sup>53, 54, 65, 67, 68, 73, 75, 90, 99, 100, 112, 115, 124</sup> A single study reporting dichotomous data showed a significant short-term effect on attitudes (OR, 1.44; 95% CI, 1.08-1.92)<sup>120</sup> (**Figure 3**).

For midterm effects of active vs control interventions (**Figure 2**), there was a significant small effect size for knowledge (SMD, 0.28; 95% CI, 0.07-0.49; 2 studies;  $I^2 = 7.5\%$ ).<sup>64,72</sup> For long-term effects, a single study showed a significant medium effect size for general stigma (SMD, 0.75; 95% CI, 0.27-1.23),<sup>65</sup> and significant small effect sizes were observed for attitudes (SMD, 0.21; 95% CI, 0.09-0.33; 2 studies;  $I^2 = 0.0\%$ )<sup>72,88</sup> and knowledge (SMD, 0.25; 95% CI, 0.08-0.42; 4 studies,  $I^2 = 0.0\%$ ),<sup>65,70,72,116</sup> along with a very small effect size for behaviors (SMD, 0.13; 95% CI, 0.03-0.22; 5 studies;  $I^2 = 0.0\%$ ).<sup>65,70,72,88,116</sup>

Moderation analyses for short-term effects of active vs control interventions for attitudes, knowledge, and general stigma found no significant subgroup differences (eTables 3-5 in [Supplement 1](#)). In contrast, significant subgroup differences were found for behaviors (eTable 6 in [Supplement 1](#)): social contact interventions had larger effect sizes than educational approaches ( $\beta = 0.15$ ; 95% CI, 0.00-0.30;  $P = .045$ ).


Short-term comparisons between active groups showed significant small effect sizes for knowledge (SMD, 0.33; 95% CI, 0.16-0.50; 9 studies;  $I^2 = 0.0\%$ )<sup>70,73,101-107</sup> and attitudes (SMD, 0.30; 95% CI, 0.02-0.57; 12 studies;  $I^2 = 30.9\%$ )<sup>55-60,73,84,96,105-107</sup> (**Figure 2**).

Moderation analyses for short-term effects between active groups on attitudes found no significant subgroup differences (eTable 7 in [Supplement 1](#)). However, significant subgroup differences were found for behaviors (eTable 8 in [Supplement 1](#)): mixed-format interventions (ie, video and nonvideo based) had smaller effect sizes than video-based formats alone ( $\beta = -0.43$ ; 95% CI, -0.83 to -0.04;  $P = .03$ ), while nonvideo-based formats had larger effect sizes than video-based formats alone ( $\beta = 1.54$ ; 95% CI, 0.04-2.67;  $P = .008$ ). Forest plots of meta-analyses of stigma-related outcomes are presented in eFigures 1 to 6 in [Supplement 1](#).

### Outcomes Associated With Help-Seeking

As shown in **Figure 2**, meta-analyses of continuous data for short-term effects of active vs control interventions showed very small significant effect sizes for help-seeking attitudes (SMD, 0.18; 95% CI, 0.09-0.28; 9 studies;  $I^2 = 0.0\%$ )<sup>67,69,75,95,97,108,112,114,124</sup> and intentions (SMD, 0.14; 95% CI, 0.07-0.21; 10 studies;  $I^2 = 0.0\%$ )<sup>61,62,75,88,99,100,108,112,115,124</sup> and a significant long-term effect size for help-seeking attitudes (SMD, 0.15; 95% CI, 0.04-0.27; 2 studies;  $I^2 = 0.0\%$ ).<sup>88,114</sup> Two studies

Figure 2. Association of Interventions With Reduced Mental Health Stigma in Young People (Continuous Outcomes)



Estimates are from 3-level multivariate meta-analyses.  $I^2_g$  indicates within-study heterogeneity;  $I^2$ , between-study heterogeneity; NA, not applicable; SMD, standardized mean difference.

reporting dichotomous data showed significant short-term effects for help-seeking attitudes (OR, 1.77; 95% CI, 1.07-2.94; 2 studies;  $I^2 = 0.0\%$ )<sup>108,120</sup> and midterm effects for formal help-seeking (OR, 1.99; 95% CI, 1.45-2.75; 2 studies;  $I^2 = 7.9\%$ )<sup>63,111</sup> and informal help-seeking (OR, 2.24; 95% CI, 1.71-2.93; 1 study)<sup>111</sup> (Figure 3). No significant effect sizes were found for short-term, midterm, or long-term effects of active vs active interventions.

Moderation analyses for short-term effects of intervention vs control conditions for help-seeking intentions found no significant subgroup differences (eTable 9 in Supplement 1). Forest plots of meta-analyses of outcomes associated with help-seeking are presented in eFigure 7 in Supplement 1.


Secondary Outcomes

No significant results were found for continuous secondary outcomes (eTable 10 in Supplement 1). A significant long-term benefit was observed between the intervention and control for helping behaviors using dichotomous data (OR, 1.43; 95% CI, 1.01-2.02; 3 studies;  $I^2 = 0.0\%$ )<sup>65,109,114</sup> (eTable 11 in Supplement 1).

Risk of Bias Across Studies

The funnel plots for short-term effects related to stigma are presented in eFigures 8 to 13 in Supplement 1. We found evidence of asymmetry in the distribution of effect sizes of active vs control interventions for attitudes, knowledge, and general stigma but not for behaviors. There was evidence of asymmetry in the distribution of effect sizes for comparisons between active groups on behaviors but not for attitudes. For help-seeking intentions, there was no evidence of asymmetry in the distribution of effect sizes for active vs control interventions (eFigure 14 in Supplement 1).

Figure 3. Association of Interventions With Reduced Mental Health Stigma in Young People (Dichotomous Outcomes)



Estimates from 3-level multivariate meta-analyses.  $I_g^2$  indicates within-study heterogeneity;  $I_s^2$ , between-study heterogeneity; NA, not applicable; OR, odds ratio.

## Discussion

We systematically reviewed 97 studies to assess the outcomes associated with interventions designed to reduce mental health stigma among young people; 74 of these studies met the criteria for 3-level multivariate meta-analyses. These studies represent a diverse sample of 43 852 young people, predominantly from developed countries. Most studies focused on college students, with educational conferences and social contact interventions being the most common approaches. Our main findings indicate that interventions yielded small to medium significant improvements in stigma-related knowledge, attitudes, behaviors, and general stigma and very small but significant improvements in help-seeking attitudes and intentions, particularly in the short term. However, the improvements tended to diminish over time, highlighting the need for sustained efforts to maintain these gains.

Our findings align with previous systematic reviews supporting the immediate benefits of mental health stigma interventions on stigma-related outcomes among young people.<sup>5,9,11,13,126</sup> The noticeable decline in positive associations over time observed in these outcomes highlights the challenges in maintaining the benefits of stigma reduction interventions, suggesting that short-term interventions may not be sufficient to produce lasting changes and that ongoing antistigma efforts, booster sessions, or follow-up activities may be necessary to sustain and enhance the initial gains.<sup>6</sup> Interestingly, social contact and nonvideo-based interventions were found to have larger effect sizes for stigma-related behaviors than educational and video-based approaches, respectively. These findings reinforce the value of incorporating direct, personal interactions with individuals with mental illness into antistigma programs compared with less personal methods.<sup>10,127-129</sup>

Our meta-analyses of a small subset of studies indicate that stigma reduction efforts may positively influence young people's willingness to seek help for mental health issues with the potential to translate into behavioral shifts.<sup>130</sup> However, effect sizes were very small, with limited evidence for long-term improvements. The limited impacts may be attributed to persistent structural barriers and cultural factors that deter readiness and ability to seek mental health support, including fears of stigma and concerns about confidentiality.<sup>21</sup> These findings highlight an important gap in antistigma interventions for promoting help-seeking behaviors. Importantly, as informal sources of help may be key in helping young people access mental health care,<sup>131</sup> more studies should consider informal help-seeking as an outcome. Complementarily, the limited significant findings related to secondary outcomes may reflect the narrower focus of current intervention designs and stress the need for more comprehensive approaches that foster long-term resilience and recovery.<sup>132</sup>

An important strength of our systematic review was the use of 3-level multivariate meta-analyses. This method provides a more nuanced understanding of within- and between-study variability; improves the precision of estimates relative to univariable meta-analyses particularly for effect sizes with higher within-study heterogeneity; and uses all available information in the primary studies, a practice often neglected in univariable meta-analyses in which researchers may be forced to opt for a particular outcome measure.<sup>18,19</sup>

The findings of this review have several implications. First, the efficacy of social contact interventions suggests that integrating lived experience narratives may be a powerful tool for reducing stigma among young people. Second, the limited long-term effects highlight the necessity of sustained and ongoing antistigma efforts that reinforce initial gains. Third, the need for more robust measures of help-seeking behaviors (ie, formal, informal) and broader psychosocial outcomes (eg, empowerment, self-efficacy) is evident. Fourth, future RCTs may benefit from using mixed-methods approaches to capture the complexity of stigma and help-seeking, providing further understanding of the factors influencing them.<sup>133</sup> Fifth, antistigma interventions were scarce from developing countries and were not found in any articles published in Spanish. As stigma is a universal phenomenon with important variations in both time and place,<sup>134</sup> more research is urgently needed in these countries where young people are particularly burdened.<sup>135</sup> Sixth, the exploration of digital interventions, which were underrepresented in this review, may offer scalable and accessible

solutions for young people worldwide, especially in challenging health contexts such as the COVID-19 pandemic. A recent study showed that COVID-19 may have had a positive influence on mental health stigma<sup>136</sup>; thus, further research on stigma and stigma interventions is needed in this new postpandemic era.

## Limitations

Important limitations of the evidence included in our review should be noted. The overall risk-of-bias assessment indicated frequent issues with masking of participants and personnel and outcome assessment, which could introduce performance and detection biases. These issues are common in behavioral intervention studies, where masking is inherently challenging.<sup>137</sup> Additionally, the unclear risk of bias in allocation concealment in more than half of the included studies points to potential selection biases that may affect the validity of the findings. Despite these limitations, the high percentage of studies with low risk of bias in random sequence generation and selective outcome reporting enhances the credibility of the reported effects. Importantly, the presence of asymmetry in the distribution of effect sizes suggests that studies with nonsignificant or negative results may be underreported, potentially inflating the observed effectiveness of interventions.

The reviewing process also had some limitations. Inclusion was limited to studies published in English or Spanish, potentially excluding important research presented in other languages. Grey literature was not included, which might have contributed to publication bias.

## Conclusions

In this systematic review and meta-analysis of 97 RCTs, we found that mental health stigma interventions for youth were associated with short-term reductions in stigma-related outcomes and improvements in help-seeking attitudes and intentions. This study provides a comprehensive overview of the current landscape of interventions aimed at reducing mental health stigma and enhancing help-seeking behaviors among young people. While the findings are encouraging, they also underscore the need for more rigorous long-term studies and the integration of multifaceted intervention strategies. By addressing these gaps, future research may contribute to the development of more effective, sustainable solutions to combat mental health stigma and support the well-being of young people globally.

## ARTICLE INFORMATION

**Accepted for Publication:** November 11, 2024.

**Published:** January 15, 2025. doi:[10.1001/jamanetworkopen.2024.54730](https://doi.org/10.1001/jamanetworkopen.2024.54730)

**Open Access:** This is an open access article distributed under the terms of the [CC-BY License](https://creativecommons.org/licenses/by/4.0/). © 2025 Crockett MA et al. *JAMA Network Open*.

**Corresponding Author:** Pablo Martínez, PhD, Douglas Mental Health University Institute, Department of Psychiatry, McGill University, 6875 Blvd LaSalle, Verdun, QC H4H 1R3, Canada ([pablo.martinezdiaz@mail.mcgill.ca](mailto:pablo.martinezdiaz@mail.mcgill.ca)).

**Author Affiliations:** Millennium Nucleus to Improve the Mental Health of Adolescents and Youths (IMHAY), Santiago, Chile (Crockett, Núñez, P. Martínez, Borghero, Campos, Langer, Carrasco, V. Martínez); Associative Research Program, Research Center on Cognitive Sciences, Facultad de Psicología, Universidad de Talca, Talca, Chile (Núñez); McGill Group for Suicide Studies, Douglas Mental Health University Institute, Department of Psychiatry, McGill University, Montreal, Quebec, Canada (P. Martínez); Millennium Institute for Research in Depression and Personality (MIDAP), Santiago, Chile (P. Martínez, V. Martínez); Center of Applied Psychology, Faculty of Psychology, Universidad de Talca, Talca, Chile (Campos); Facultad de Psicología y Humanidades, Universidad San Sebastián, Valdivia, Chile (Langer); Facultad de Medicina, Universidad Austral de Chile, Valdivia, Chile (Carrasco); Centro de Medicina Reproductiva y Desarrollo Integral del Adolescente (CEMERA), Facultad de Medicina, Universidad de Chile, Santiago, Chile (V. Martínez).

**Author Contributions:** Dr P. Martínez had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

**Concept and design:** Núñez, P. Martínez, Borghero, Campos, V. Martínez.

**Acquisition, analysis, or interpretation of data:** Crockett, Núñez, P. Martínez, Borghero, Langer, Carrasco, V. Martínez.

**Drafting of the manuscript:** Crockett, Núñez, P. Martínez, Borghero, Carrasco, V. Martínez.

**Critical review of the manuscript for important intellectual content:** Crockett, Núñez, Campos, Langer, V. Martínez.

**Statistical analysis:** Crockett, P. Martínez.

**Obtained funding:** Carrasco, V. Martínez.

**Administrative, technical, or material support:** Crockett, Núñez, P. Martínez, Borghero, Campos, V. Martínez.

**Supervision:** Núñez, P. Martínez, Langer, V. Martínez.

**Conflict of Interest Disclosures:** None reported.

**Funding/Support:** This study was funded by Millennium Science Initiative Program grants NCS2021\_081 and ICS13\_005 and National Fund for Scientific and Technological Development grants 1221230 and 1210093 from the National Agency of Research and Development.

**Role of the Funder/Sponsor:** The funder had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

**Data Sharing Statement:** See [Supplement 2](#).

## REFERENCES

1. Sheehan L, Nieweglowski K, Corrigan PW. Structures and types of stigma. In: Gaebel W, Rössler W, Sartorius N, eds. *The Stigma of Mental Illness—End of the Story?* Springer International Publishing; 2017:43-66. doi:[10.1007/978-3-319-27839-1\\_3](#)
2. Clement S, Schauman O, Graham T, et al. What is the impact of mental health-related stigma on help-seeking? a systematic review of quantitative and qualitative studies. *Psychol Med*. 2015;45(1):11-27. doi:[10.1017/S0033291714000129](#)
3. Ebert DD, Mortier P, Kaehlke F, et al; WHO World Mental Health-International College Student Initiative Collaborators. Barriers of mental health treatment utilization among first-year college students: first cross-national results from the WHO World Mental Health International College Student Initiative. *Int J Methods Psychiatr Res*. 2019;28(2):e1782. doi:[10.1002/mpr.1782](#)
4. World Health Organization. *World Mental Health Report: Transforming Mental Health for All*. World Health Organization; 2022.
5. Thornicroft G, Sunkel C, Alikhon Aliev A, et al. The Lancet Commission on ending stigma and discrimination in mental health. *Lancet*. 2022;400(10361):1438-1480. doi:[10.1016/S0140-6736\(22\)01470-2](#)
6. Clayborne ZM, Varin M, Colman I. Systematic review and meta-analysis: adolescent depression and long-term psychosocial outcomes. *J Am Acad Child Adolesc Psychiatry*. 2019;58(1):72-79. doi:[10.1016/j.jaac.2018.07.896](#)
7. Costello EJ, Maughan B. Annual research review: optimal outcomes of child and adolescent mental illness. *J Child Psychol Psychiatry*. 2015;56(3):324-341. doi:[10.1111/jcpp.12371](#)
8. Erskine HE, Norman RE, Ferrari AJ, et al. Long-term outcomes of attention-deficit/hyperactivity disorder and conduct disorder: a systematic review and meta-analysis. *J Am Acad Child Adolesc Psychiatry*. 2016;55(10):841-850. doi:[10.1016/j.jaac.2016.06.016](#)
9. Parry SJ, Brohan E, Gronholm PC, Thornicroft G. Interventions to reduce mental illness stigma and discrimination at the person-level for individuals and small groups. In: Vogel DL, Wade NG, eds. *The Cambridge Handbook of Stigma and Mental Health*. Cambridge University Press; 2022:369-390. doi:[10.1017/9781108920995.022](#)
10. Corrigan PW, Morris SB, Michaels PJ, Rafacz JD, Rüsch N. Challenging the public stigma of mental illness: a meta-analysis of outcome studies. *Psychiatr Serv*. 2012;63(10):963-973. doi:[10.1176/appi.ps.201100529](#)
11. Mellor C. School-based interventions targeting stigma of mental illness: systematic review. *Psychiatr Bull* (2014). 2014;38(4):164-171. doi:[10.1192/pb.bp.112.041723](#)
12. Schachter HM, Girardi A, Ly M, et al. Effects of school-based interventions on mental health stigmatization: a systematic review. *Child Adolesc Psychiatry Ment Health*. 2008;2(1):18. doi:[10.1186/1753-2000-2-18](#)
13. Waqas A, Malik S, Fida A, et al. Interventions to reduce stigma related to mental illnesses in educational institutes: a systematic review. *Psychiatr Q*. 2020;91(3):887-903. doi:[10.1007/s11126-020-09751-4](#)

14. Heim E, Henderson C, Kohrt BA, Koschorke M, Milenova M, Thornicroft G. Reducing mental health-related stigma among medical and nursing students in low- and middle-income countries: a systematic review. *Epidemiol Psychiatr Sci*. 2019;29:e28. doi:[10.1017/S2045796019000167](https://doi.org/10.1017/S2045796019000167)
15. Yamaguchi S, Wu SI, Biswas M, et al. Effects of short-term interventions to reduce mental health-related stigma in university or college students: a systematic review. *J Nerv Ment Dis*. 2013;201(6):490-503. doi:[10.1097/NMD.0b013e31829480df](https://doi.org/10.1097/NMD.0b013e31829480df)
16. Song N, Hugh-Jones S, West RM, Pickavance J, Mir G. The effectiveness of anti-stigma interventions for reducing mental health stigma in young people: a systematic review and meta-analysis. *Glob Ment Health (Camb)*. 2023;10:e39. doi:[10.1017/gmh.2023.34](https://doi.org/10.1017/gmh.2023.34)
17. Gaiha SM, Salisbury TT, Usmani S, Koschorke M, Raman U, Petticrew M. Effectiveness of arts interventions to reduce mental-health-related stigma among youth: a systematic review and meta-analysis. *BMC Psychiatry*. 2021;21(1):364. doi:[10.1186/s12888-021-03350-8](https://doi.org/10.1186/s12888-021-03350-8)
18. Riley RD. Multivariate meta-analysis: the effect of ignoring within-study correlation. *J R Stat Soc Ser A Stat Soc*. 2009;172(4):789-811. doi:[10.1111/j.1467-985X.2008.00593.x](https://doi.org/10.1111/j.1467-985X.2008.00593.x)
19. Wei Y, Higgins JP. Estimating within-study covariances in multivariate meta-analysis with multiple outcomes. *Stat Med*. 2013;32(7):1191-1205. doi:[10.1002/sim.5679](https://doi.org/10.1002/sim.5679)
20. Núñez D, Martínez P, Borghero F, Campos S, Martínez V. Interventions to reduce stigma towards mental disorders in young people: protocol for a systematic review and meta-analysis. *BMJ Open*. 2021;11(11):e045726. doi:[10.1136/bmjopen-2020-045726](https://doi.org/10.1136/bmjopen-2020-045726)
21. Rickwood D, Thomas K. Conceptual measurement framework for help-seeking for mental health problems. *Psychol Res Behav Manag*. 2012;5:173-183. doi:[10.2147/PRBM.S38707](https://doi.org/10.2147/PRBM.S38707)
22. Higgins JPT, Altman DG. Assessing risk of bias in included studies. In: Higgins JPT, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions*. Wiley; 2008:187-241. doi:[10.1002/9780470712184.ch8](https://doi.org/10.1002/9780470712184.ch8)
23. Higgins J, Li T, Deeks J. Choosing effect measures and computing estimates of effect. In: Higgins JPT, Thomas J, Chandler J, et al, eds. *Cochrane Handbook for Systematic Reviews of Interventions Version 63 (updated February 2022)*. Cochrane; 2022:chap 6.
24. Deeks JJ, Higgins JP, Altman DG; Cochrane Statistical Methods Group. Analysing data and undertaking meta-analyses. In: Higgins JPT, Thomas J, Chandler J, et al, eds. *Cochrane Handbook for Systematic Reviews of Interventions*. Wiley; 2019:241-284. doi:[10.1002/978119536604.ch10](https://doi.org/10.1002/978119536604.ch10)
25. Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J Stat Softw*. 2010;36(3). doi:[10.18637/jss.v036.i03](https://doi.org/10.18637/jss.v036.i03)
26. Amsalem D, Jankowski SE, Pagdon S, et al. "It is hard to be a woman with schizophrenia": randomized controlled trial of a brief video intervention to reduce public stigma in young adults. *J Clin Psychiatry*. 2022;84(1):22m14534. doi:[10.4088/JCP.22m14534](https://doi.org/10.4088/JCP.22m14534)
27. Boucher LA, Campbell DG. An examination of the impact of a biological anti-stigma message for depression on college students. *J Coll Stud Psychother*. 2014;28(1):74-81. doi:[10.1080/87568225.2014.854686](https://doi.org/10.1080/87568225.2014.854686)
28. Campbell M, Shryane N, Byrne R, Morrison AP. A mental health promotion approach to reducing discrimination about psychosis in teenagers. *Psychosis*. 2011;3(1):41-51. doi:[10.1080/17522431003735529](https://doi.org/10.1080/17522431003735529)
29. Cheetham A, Sandral E, Lubman DI. Outcomes of reducing stigma towards alcohol misuse during adolescence: results of a randomized controlled trial of the MAKINGtheLINK intervention. *Child Adolesc Psychiatry Ment Health*. 2020;14(1):11. doi:[10.1186/s13034-020-00317-7](https://doi.org/10.1186/s13034-020-00317-7)
30. Chung KF. Changing the attitudes of Hong Kong medical students toward people with mental illness. *J Nerv Ment Dis*. 2005;193(11):766-768. doi:[10.1097/01.nmd.0000185871.20390.c6](https://doi.org/10.1097/01.nmd.0000185871.20390.c6)
31. Chung KF, Chan JH. Can a less pejorative Chinese translation for schizophrenia reduce stigma? a study of adolescents' attitudes toward people with schizophrenia. *Psychiatry Clin Neurosci*. 2004;58(5):507-515. doi:[10.1111/j.1440-1819.2004.01293.x](https://doi.org/10.1111/j.1440-1819.2004.01293.x)
32. Conceição V, Rothes I, Gusmão R. The effects of a video-based randomized controlled trial intervention on depression stigma and help-seeking attitudes in university students. *Psychiatry Res*. 2022;308:114356. doi:[10.1016/j.psychres.2021.114356](https://doi.org/10.1016/j.psychres.2021.114356)
33. Corrigan PW, Rafacz JD, Hautamaki J, et al. Changing stigmatizing perceptions and recollections about mental illness: the effects of NAMI's in Our Own Voice. *Community Ment Health J*. 2010;46(5):517-522. doi:[10.1007/s10597-009-9287-3](https://doi.org/10.1007/s10597-009-9287-3)
34. Economou M, Louki E, Peppou LE, Gramandani C, Yotis L, Stefanis CN. Fighting psychiatric stigma in the classroom: the impact of an educational intervention on secondary school students' attitudes to schizophrenia. *Int J Soc Psychiatry*. 2012;58(5):544-551. doi:[10.1177/0020764011413678](https://doi.org/10.1177/0020764011413678)

35. Economou M, Peppou LE, Geroulanou K, et al. The influence of an anti-stigma intervention on adolescents' attitudes to schizophrenia: a mixed methodology approach. *Child Adolesc Ment Health*. 2014;19(1):16-23. doi:10.1111/j.1475-3588.2012.00669.x
36. Esanu A, Chihai J, Esanu D. Educational intervention for adolescents in order to decrease stigma towards people with mental disorders: cluster randomized controlled trial. *Eur Neuropsychopharmacol*. 2016;26:S739. doi:10.1016/S0924-977X(16)31894-6
37. Han J, Batterham PJ, Calear AL, Wu Y, Xue J, van Spijker BAJ. Development and pilot evaluation of an online psychoeducational program for suicide prevention among university students: a randomised controlled trial. *Internet Interv*. 2017;12:111-120. doi:10.1016/j.invent.2017.11.002
38. Koike S, Yamaguchi S, Ojio Y, et al. A randomised controlled trial of repeated filmed social contact on reducing mental illness-related stigma in young adults. *Epidemiol Psychiatr Sci*. 2018;27(2):199-208. doi:10.1017/S2045796016001050
39. Lannin DG, Guyll M, Vogel DL, Madon S. Reducing the stigma associated with seeking psychotherapy through self-affirmation. *J Couns Psychol*. 2013;60(4):508-519. doi:10.1037/a0033789
40. Lincoln TM, Arens E, Berger C, Rief W. Can antistigma campaigns be improved? a test of the impact of biogenetic vs psychosocial causal explanations on implicit and explicit attitudes to schizophrenia. *Schizophr Bull*. 2008;34(5):984-994. doi:10.1093/schbul/sbm131
41. Mann CE, Himelein MJ. Putting the person back into psychopathology: an intervention to reduce mental illness stigma in the classroom. *Soc Psychiatry Psychiatr Epidemiol*. 2008;43(7):545-551. doi:10.1007/s00127-008-0324-2
42. Mulfinger N, Müller S, Böge I, et al. Honest, Open, Proud for adolescents with mental illness: pilot randomized controlled trial. *J Child Psychol Psychiatry*. 2018;59(6):684-691. doi:10.1111/jcpp.12853
43. O'Kearney R, Gibson M, Christensen H, Griffiths KM. Effects of a cognitive-behavioural internet program on depression, vulnerability to depression and stigma in adolescent males: a school-based controlled trial. *Cogn Behav Ther*. 2006;35(1):43-54. doi:10.1080/16506070500303456
44. O'Mara L, Akhtar-Danesh N, Browne G, et al. Does youth net decrease mental illness stigma in high school students? *Can J Commun Ment Health*. 2013;32(4):9-22. doi:10.7870/cjcmh-2013-023
45. Primbs R, Feldmann L, Iglhaut L, Allgaier AK, Schulte-Körne G, Greimel E. Evaluation of an information booklet for adolescents on depression: evidence from a randomized controlled study. *Child Adolesc Psychiatry Ment Health*. 2023;17(1):65. doi:10.1186/s13034-023-00614-x
46. Queroue M, Pouymayou A, Pereira E, Tzourio C, González-Caballero JL, Montagni I. An interactive video increasing French students' mental health literacy: a mixed-methods randomized controlled pilot study. *Health Promot Int*. 2023;38(4):daab202. doi:10.1093/heapro/daab202
47. Rainone N, Oodal R, Niederdeppe J. The (surprising) impact of televised antidepressant direct-to-consumer advertising on the stigmatization of mental illness. *Community Ment Health J*. 2018;54(3):267-275. doi:10.1007/s10597-017-0164-1
48. Saif N, Woodbury M, Hodes J, et al. Effectiveness of online Alzheimer's and brain health education for high school and college students: a randomized controlled trial. *Alzheimers Dement*. 2019;15(75 pt 23):P1202-P1204. doi:10.1016/j.jalz.2019.06.3632
49. Tan ASY, Mankiewicz PD. Does a brief contact-based video intervention reduce public mental health stigma associated with psychosis in a collectivistic society? a randomised control study. *J Ment Health Train Educ Pract*. 2024;19(1):21-35. doi:10.1108/JMHTEP-05-2023-0053
50. Taylor-Rodgers E, Batterham PJ. Evaluation of an online psychoeducation intervention to promote mental health help seeking attitudes and intentions among young adults: randomised controlled trial. *J Affect Disord*. 2014;168:65-71. doi:10.1016/j.jad.2014.06.047
51. Greif Green J, Levine RS, Oblath R, Corriveau KH, Holt MK, Albright G. Pilot evaluation of preservice teacher training to improve preparedness and confidence to address student mental health. *Evid Based Pract Child Adolesc Ment Health*. 2020;5(1):42-52. doi:10.1080/23794925.2020.1727793
52. Mankiewicz PD, Tan AS. Reducing discrimination against service users with psychosis through brief contact-based video interventions: a randomised controlled trial. *Br J Healthc Manag*. 2024;30(1):14-24. doi:10.12968/bjhc.2023.0074
53. Pos K, Franke N, Smit F, et al. Cognitive behavioral therapy for social activation in recent-onset psychosis: randomized controlled trial. *J Consult Clin Psychol*. 2019;87(2):151-160. doi:10.1037/ccp0000362

54. Yuen ASY, Mak WWS. The effects of immersive virtual reality in reducing public stigma of mental illness in the university population of Hong Kong: randomized controlled trial. *J Med Internet Res*. 2021;23(7):e23683. doi:10.2196/23683
55. Finkelstein J, Lapshin O, Wasserman E. Randomized study of different anti-stigma media. *Patient Educ Couns*. 2008;71(2):204-214. doi:10.1016/j.pec.2008.01.002
56. Brown SA. Implementing a brief hallucination simulation as a mental illness stigma reduction strategy. *Community Ment Health J*. 2010;46(5):500-504. doi:10.1007/s10597-009-9229-0
57. Fernandez A, Tan KA, Knaak S, Chew BH, Ghazali SS. Effects of brief psychoeducational program on stigma in Malaysian pre-clinical medical students: a randomized controlled trial. *Acad Psychiatry*. 2016;40(6):905-911. doi:10.1007/s40596-016-0592-1
58. King CA, Eisenberg D, Zheng K, et al. Online suicide risk screening and intervention with college students: a pilot randomized controlled trial. *J Consult Clin Psychol*. 2015;83(3):630-636. doi:10.1037/a0038805
59. Masuda A, Hayes SC, Fletcher LB, et al. Impact of acceptance and commitment therapy versus education on stigma toward people with psychological disorders. *Behav Res Ther*. 2007;45(11):2764-2772. doi:10.1016/j.brat.2007.05.008
60. Winkler P, Janoušková M, Kožený J, et al. Short video interventions to reduce mental health stigma: a multi-centre randomised controlled trial in nursing high schools. *Soc Psychiatry Psychiatr Epidemiol*. 2017;52(12):1549-1557. doi:10.1007/s00127-017-1449-y
61. Han DY, Chen SH, Hwang KK, Wei HL. Effects of psychoeducation for depression on help-seeking willingness: biological attribution versus destigmatization. *Psychiatry Clin Neurosci*. 2006;60(6):662-668. doi:10.1111/j.1440-1819.2006.01580.x
62. Kruger E, Pitts SC, Denenny D, DeLuca JS, Schiffman J. Efficacy of contact intervention videos on college students' intentions toward mental health help-seeking. *J Am Coll Health*. 2022;70(7):2033-2041. doi:10.1080/07448481.2020.1842417
63. da Conceição V, Mesquita E, Gusmão R. Effects of a stigma reduction intervention on help-seeking behaviors in university students: a 2019-2021 randomized controlled trial. *Psychiatry Res*. 2024;331:115673. doi:10.1016/j.psychres.2023.115673
64. Campos L, Dias P, Duarte A, Veiga E, Dias CC, Palha F. Is it possible to "find space for mental health" in young people? effectiveness of a school-based mental health literacy promotion program. *Int J Environ Res Public Health*. 2018;15(7):1426. doi:10.3390/ijerph15071426
65. Dzemaili S, Pasquier J, Oulevey Bachmann A, Mohler-Kuo M. The effectiveness of mental health first aid training among undergraduate students in Switzerland: a randomized control trial. *Int J Environ Res Public Health*. 2023;20(2):1303. doi:10.3390/ijerph20021303
66. Mori R, Uchino T, Mizuno M, Yamaguchi T, Katagiri N, Nemoto T. Effectiveness of a comprehensive mental health literacy educational programme for junior high school students: a randomised controlled trial examining changes in their knowledge, attitudes, and behaviour. *J Pers Med*. 2022;12(8):1281. doi:10.3390/jpm12081281
67. Simkiss NJ, Gray NS, Kemp AH, Dunne C, Snowden RJ. A randomised controlled trial evaluating the Guide Cymru mental health literacy intervention programme in year 9 (age 13-14) school pupils in Wales. *BMC Public Health*. 2023;23(1):1062. doi:10.1186/s12889-023-15922-2
68. Tay JL, Goh YSS, Sim K, Klainin-Yobas P. Impact of the HOPE intervention on mental health literacy, psychological well-being and stress levels amongst university undergraduates: a randomised controlled trial. *Int J Environ Res Public Health*. 2022;19(15):9772. doi:10.3390/ijerph19159772
69. Oshima F, Mandy W, Seto M, et al. Cognitive behavior therapy for autistic adolescents, awareness and care for my autistic traits program: a multicenter randomized controlled trial. *BMC Psychiatry*. 2023;23(1):661. doi:10.1186/s12888-023-05075-2
70. Yamaguchi S, Ojio Y, Ando S, et al. Long-term effects of filmed social contact or internet-based self-study on mental health-related stigma: a 2-year follow-up of a randomised controlled trial. *Soc Psychiatry Psychiatr Epidemiol*. 2019;54(1):33-42. doi:10.1007/s00127-018-1609-8
71. Brown SA. The effects of direct-to-consumer-advertising on mental illness beliefs and stigma. *Community Ment Health J*. 2017;53(5):534-541. doi:10.1007/s10597-017-0121-z
72. Casañas R, Castellvi P, Gil JJ, et al. The effectiveness of a "EspaiJove.net" - a school-based intervention programme in increasing mental health knowledge, help seeking and reducing stigma attitudes in the adolescent population: a cluster randomised controlled trial. *BMC Public Health*. 2022;22(1):2425. doi:10.1186/s12889-022-14558-y

73. Chung EY, Tse TT. Effect of human library intervention on mental health literacy: a multigroup pretest-posttest study. *BMC Psychiatry*. 2022;22(1):73. doi:[10.1186/s12888-022-03725-5](https://doi.org/10.1186/s12888-022-03725-5)
74. Farina N, Hughes LJ, Jones E, et al. The effect of a dementia awareness class on changing dementia attitudes in adolescents. *BMC Geriatr*. 2020;20(1):188. doi:[10.1186/s12877-020-01589-6](https://doi.org/10.1186/s12877-020-01589-6)
75. Gulliver A, Griffiths KM, Christensen H, et al. Internet-based interventions to promote mental health help-seeking in elite athletes: an exploratory randomized controlled trial. *J Med Internet Res*. 2012;14(3):e69. doi:[10.2196/jmir.1864](https://doi.org/10.2196/jmir.1864)
76. Lipson SK, Speer N, Brunwasser S, Hahn E, Eisenberg D. Gatekeeper training and access to mental health care at universities and colleges. *J Adolesc Health*. 2014;55(5):612-619. doi:[10.1016/j.jadohealth.2014.05.009](https://doi.org/10.1016/j.jadohealth.2014.05.009)
77. Milin R, Kutcher S, Lewis SP, et al. Impact of a mental health curriculum on knowledge and stigma among high school students: a randomized controlled trial. *J Am Acad Child Adolesc Psychiatry*. 2016;55(5):383-391.e1. doi:[10.1016/j.jaac.2016.02.018](https://doi.org/10.1016/j.jaac.2016.02.018)
78. Oduguwa AO, Adedokun B, Omigbodun OO. Effect of a mental health training programme on Nigerian school pupils' perceptions of mental illness. *Child Adolesc Psychiatry Ment Health*. 2017;11(1):19. doi:[10.1186/s13034-017-0157-4](https://doi.org/10.1186/s13034-017-0157-4)
79. Penn DL, Chamberlin C, Mueser KT. The effects of a documentary film about schizophrenia on psychiatric stigma. *Schizophr Bull*. 2003;29(2):383-391. doi:[10.1093/oxfordjournals.schbul.a007012](https://doi.org/10.1093/oxfordjournals.schbul.a007012)
80. Ranson NJ, Byrne MK. Promoting peer acceptance of females with higher-functioning autism in a mainstream education setting: a replication and extension of the effects of an autism anti-stigma program. *J Autism Dev Disord*. 2014;44(11):2778-2796. doi:[10.1007/s10803-014-2139-1](https://doi.org/10.1007/s10803-014-2139-1)
81. Staniland JJ, Byrne MK. The effects of a multi-component higher-functioning autism anti-stigma program on adolescent boys. *J Autism Dev Disord*. 2013;43(12):2816-2829. doi:[10.1007/s10803-013-1829-4](https://doi.org/10.1007/s10803-013-1829-4)
82. Wood AL, Wahl OF. Evaluating the effectiveness of a consumer-provided mental health recovery education presentation. *Psychiatr Rehabil J*. 2006;30(1):46-53. doi:[10.2975/30.2006.46.53](https://doi.org/10.2975/30.2006.46.53)
83. Pinto-Foltz MD, Logsdon MC, Myers JA. Feasibility, acceptability, and initial efficacy of a knowledge-contact program to reduce mental illness stigma and improve mental health literacy in adolescents. *Soc Sci Med*. 2011;72(12):2011-2019. doi:[10.1016/j.socscimed.2011.04.006](https://doi.org/10.1016/j.socscimed.2011.04.006)
84. Brown SA, Evans Y, Espenschied K, O'Connor M. An examination of two brief stigma reduction strategies: filmed personal contact and hallucination simulations. *Community Ment Health J*. 2010;46(5):494-499. doi:[10.1007/s10597-010-9309-1](https://doi.org/10.1007/s10597-010-9309-1)
85. Giannakopoulos G, Assimopoulos H, Petanidou D, Tzavara C, Kolaitis G, Tsiantis J. Effectiveness of a school-based intervention for enhancing adolescents' positive attitudes towards people with mental illness. *Ment Illn*. 2012;4(2):e16. doi:[10.4081/mi.2012.e16](https://doi.org/10.4081/mi.2012.e16)
86. Grandón P, Saldivia S, Cova F, et al. Effectiveness of an intervention to reduce stigma towards people with a severe mental disorder diagnosis in primary health care personnel: Programme Igual-Mente. *Psychiatry Res*. 2021;305:114259. doi:[10.1016/j.psychres.2021.114259](https://doi.org/10.1016/j.psychres.2021.114259)
87. Han DY, Chen SH. Reducing the stigma of depression through neurobiology-based psychoeducation: a randomized controlled trial. *Psychiatry Clin Neurosci*. 2014;68(9):666-673. doi:[10.1111/pcn.12174](https://doi.org/10.1111/pcn.12174)
88. Hart LM, Morgan AJ, Rossetto A, Kelly CM, Mackinnon A, Jorm AF. Helping adolescents to better support their peers with a mental health problem: a cluster-randomised crossover trial of teen Mental Health First Aid. *Aust N Z J Psychiatry*. 2018;52(7):638-651. doi:[10.1177/0004867417753552](https://doi.org/10.1177/0004867417753552)
89. Kerby J, Calton T, Dimambro B, Flood C, Glazebrook C. Anti-stigma films and medical students' attitudes towards mental illness and psychiatry: randomised controlled trial. *Psychiatr Bull*. 2008;32(9):345-349. doi:[10.1192/pb.bp.107.017152](https://doi.org/10.1192/pb.bp.107.017152)
90. Rubio-Valera M, Aznar-Lou I, Vives-Collet M, Fernández A, Gil-Girbau M, Serrano-Blanco A. Reducing the mental health-related stigma of social work students: a cluster RCT. *Res Soc Work Pract*. 2018;28(2):164-172. doi:[10.1177/1049731516641492](https://doi.org/10.1177/1049731516641492)
91. Bruins J, Bartels-Velthuis AA, van Weeghel J, Helmus K, Pijnenborg GHM. Letter to the editors: reducing stigmatizing attitudes in high school adolescents—a cluster RCT on the effectiveness of a schizophrenia documentary. *Schizophr Res*. 2018;202:408-409. doi:[10.1016/j.schres.2018.06.055](https://doi.org/10.1016/j.schres.2018.06.055)
92. Rahman A, Mubbashar MH, Gater R, Goldberg D. Randomised trial of impact of school mental-health programme in rural Rawalpindi, Pakistan. *Lancet*. 1998;352(9133):1022-1025. doi:[10.1016/S0140-6736\(98\)02381-2](https://doi.org/10.1016/S0140-6736(98)02381-2)
93. Vila-Badia R, Martínez-Zambrano F, Arenas O, et al. Effectiveness of an intervention for reducing social stigma towards mental illness in adolescents. *World J Psychiatry*. 2016;6(2):239-247. doi:[10.5498/wjp.v6.i2.239](https://doi.org/10.5498/wjp.v6.i2.239)

94. Davies EB, Beever E, Glazebrook C. A pilot randomised controlled study of the Mental Health First Aid eLearning course with UK medical students. *BMC Med Educ*. 2018;18(1):45. doi:10.1186/s12909-018-1154-x
95. Kirschner B, Goetzl M, Curtin L. Mental health stigma among college students: test of an interactive online intervention. *J Am Coll Health*. 2022;70(6):1831-1838. doi:10.1080/07448481.2020.1826492
96. Rusch LC, Kanter JW, Brondino MJ. A comparison of contextual and biomedical models of stigma reduction for depression with a nonclinical undergraduate sample. *J Nerv Ment Dis*. 2009;197(2):104-110. doi:10.1097/NMD.0b013e318192416f
97. Sharp W, Hargrove DS, Johnson L, Deal WP. Mental health education: an evaluation of a classroom based strategy to modify help seeking for mental health problems. *J Coll Student Dev*. 2006;47(4):419-438. doi:10.1353/csd.2006.0051
98. Amsalem D, Halloran J, Penque B, Celentano J, Martin A. Effect of a brief social contact video on transphobia and depression-related stigma among adolescents: a randomized clinical trial. *JAMA Netw Open*. 2022;5(2):e220376. doi:10.1001/jamanetworkopen.2022.0376
99. Amsalem D, Jankowski SE, Pagdon S, et al. Selfie videos to reduce stigma and increase treatment seeking among youths: two noninferiority randomized controlled trials. *Psychiatr Serv*. 2023;74(3):229-236. doi:10.1176/appi.ps.20220168
100. Amsalem D, Martin A. Reducing depression-related stigma and increasing treatment seeking among adolescents: randomized controlled trial of a brief video intervention. *J Child Psychol Psychiatry*. 2022;63(2):210-217. doi:10.1111/jcpp.13427
101. Chisholm K, Patterson P, Torgerson C, Turner E, Jenkinson D, Birchwood M. Impact of contact on adolescents' mental health literacy and stigma: the SchoolSpace cluster randomised controlled trial. *BMJ Open*. 2016;6(2):e009435. doi:10.1136/bmjopen-2015-009435
102. Han M, Wang Y, Zhang Y, et al. A multicomponent digital intervention to promote help-seeking for mental health problems and suicide in sexual and gender diverse young adults: a randomized controlled trial. *PLoS Med*. 2023;20(3):e1004197. doi:10.1371/journal.pmed.1004197
103. Ojio Y, Yamaguchi S, Ohta K, Ando S, Koike S. Effects of biomedical messages and expert-recommended messages on reducing mental health-related stigma: a randomised controlled trial. *Epidemiol Psychiatr Sci*. 2019;29:e74. doi:10.1017/S2045796019000714
104. Perry Y, Petrie K, Buckley H, et al. Effects of a classroom-based educational resource on adolescent mental health literacy: a cluster randomized controlled trial. *J Adolesc*. 2014;37(7):1143-1151. doi:10.1016/j.adolescence.2014.08.001
105. Chan JYN, Mak WWS, Law LSC. Combining education and video-based contact to reduce stigma of mental illness: "The Same or Not the Same" anti-stigma program for secondary schools in Hong Kong. *Soc Sci Med*. 2009;68(8):1521-1526. doi:10.1016/j.socscimed.2009.02.016
106. Corrigan PW, Larson J, Sells M, Niessen N, Watson AC. Will filmed presentations of education and contact diminish mental illness stigma? *Community Ment Health J*. 2007;43(2):171-181. doi:10.1007/s10597-006-9061-8
107. Norman RMG, Li Y, Sorrentino R, Hampson E, Ye Y. The differential effects of a focus on symptoms versus recovery in reducing stigma of schizophrenia. *Soc Psychiatry Psychiatr Epidemiol*. 2017;52(11):1385-1394. doi:10.1007/s00127-017-1429-2
108. Costin DL, Mackinnon AJ, Griffiths KM, et al. Health e-cards as a means of encouraging help seeking for depression among young adults: randomized controlled trial. *J Med Internet Res*. 2009;11(4):e42. doi:10.2196/jmir.1294
109. Hart LM, Morgan AJ, Rossetto A, et al. Teen Mental Health First Aid: 12-month outcomes from a cluster crossover randomized controlled trial evaluation of a universal program to help adolescents better support peers with a mental health problem. *BMC Public Health*. 2022;22(1):1159. doi:10.1186/s12889-022-13554-6
110. Howard KA, Griffiths KM, McKetin R, Ma J. Can a brief biologically-based psychoeducational intervention reduce stigma and increase help-seeking intentions for depression in young people? a randomised controlled trial. *J Child Adolesc Ment Health*. 2018;30(1):27-39. doi:10.2989/17280583.2018.1467323
111. Jorm AF, Kitchener BA, Sawyer MG, Scales H, Cvetkovski S. Mental Health First Aid training for high school teachers: a cluster randomized trial. *BMC Psychiatry*. 2010;10(1):51. doi:10.1186/1471-244X-10-51
112. Kosyluk KA, Al-Khouja M, Bink A, et al. Challenging the stigma of mental illness among college students. *J Adolesc Health*. 2016;59(3):325-331. doi:10.1016/j.jadohealth.2016.05.005
113. Link BG, DuPont-Reyes MJ, Barkin K, Villatoro AP, Phelan JC, Painter K. A school-based intervention for mental illness stigma: a cluster randomized trial. *Pediatrics*. 2020;145(6):e20190780. doi:10.1542/peds.2019-0780

114. Lubman DI, Cheetham A, Sandral E, et al. Twelve-month outcomes of MAKINGtheLINK: a cluster randomized controlled trial of a school-based program to facilitate help-seeking for substance use and mental health problems. *EClinicalMedicine*. 2020;18:100225. doi:[10.1016/j.eclinm.2019.11.018](https://doi.org/10.1016/j.eclinm.2019.11.018)
115. Martin A, Calhoun A, Pérez J, Amsalem D. Destigmatizing perceptions about Black adolescent depression: randomized controlled trial of brief social contact-based video interventions. *J Child Psychol Psychiatry*. 2022;63(11):1270-1278. doi:[10.1111/jcpp.13570](https://doi.org/10.1111/jcpp.13570)
116. Morgan AJ, Fischer JAA, Hart LM, et al. Does Mental Health First Aid training improve the mental health of aid recipients? the training for parents of teenagers randomised controlled trial. *BMC Psychiatry*. 2019;19(1):99. doi:[10.1186/s12888-019-2085-8](https://doi.org/10.1186/s12888-019-2085-8)
117. Morrison AP, Birchwood M, Pyle M, et al. Impact of cognitive therapy on internalised stigma in people with at-risk mental states. *Br J Psychiatry*. 2013;203(2):140-145. doi:[10.1192/bjp.bp.112.123703](https://doi.org/10.1192/bjp.bp.112.123703)
118. Papish A, Kassam A, Modgill G, Vaz G, Zanussi L, Patten S. Reducing the stigma of mental illness in undergraduate medical education: a randomized controlled trial. *BMC Med Educ*. 2013;13(1):141. doi:[10.1186/1472-6920-13-141](https://doi.org/10.1186/1472-6920-13-141)
119. Patten SB, Remillard A, Phillips L, et al. Effectiveness of contact-based education for reducing mental illness-related stigma in pharmacy students. *BMC Med Educ*. 2012;12(1):120. doi:[10.1186/1472-6920-12-120](https://doi.org/10.1186/1472-6920-12-120)
120. Reavley NJ, McCann TV, Cvetkovski S, Jorm AF. A multifaceted intervention to improve mental health literacy in students of a multicampus university: a cluster randomised trial. *Soc Psychiatry Psychiatr Epidemiol*. 2014;49(10):1655-1666. doi:[10.1007/s00127-014-0880-6](https://doi.org/10.1007/s00127-014-0880-6)
121. Sebastian J, Richards D. Changing stigmatizing attitudes to mental health via education and contact with embodied conversational agents. *Comput Human Behav*. 2017;73:479-488. doi:[10.1016/j.chb.2017.03.071](https://doi.org/10.1016/j.chb.2017.03.071)
122. Stanley IH, Hom MA, Joiner TE. Modifying mental health help-seeking stigma among undergraduates with untreated psychiatric disorders: a pilot randomized trial of a novel cognitive bias modification intervention. *Behav Res Ther*. 2018;103:33-42. doi:[10.1016/j.brat.2018.01.008](https://doi.org/10.1016/j.brat.2018.01.008)
123. Swartz K, Musci RJ, Beaudry MB, et al. School-based curriculum to improve depression literacy among US secondary school students: a randomized effectiveness trial. *Am J Public Health*. 2017;107(12):1970-1976. doi:[10.2105/AJPH.2017.304088](https://doi.org/10.2105/AJPH.2017.304088)
124. Tuijnman A, Kleinjan M, Olthof M, Hoogendoorn E, Granic I, Engels RC. A game-based school program for mental health literacy and stigma on depression (Moving Stories): cluster randomized controlled trial. *JMIR Ment Health*. 2022;9(8):e26615. doi:[10.2196/26615](https://doi.org/10.2196/26615)
125. Wright A, McGorry PD, Harris MG, Jorm AF, Pennell K. Development and evaluation of a youth mental health community awareness campaign—the Compass Strategy. *BMC Public Health*. 2006;6(1):215. doi:[10.1186/1471-2458-6-215](https://doi.org/10.1186/1471-2458-6-215)
126. Ma KKY, Anderson JK, Burn AM. Review: school-based interventions to improve mental health literacy and reduce mental health stigma—a systematic review. *Child Adolesc Ment Health*. 2023;28(2):230-240. doi:[10.1111/camh.12543](https://doi.org/10.1111/camh.12543)
127. Gronholm PC, Henderson C, Deb T, Thornicroft G. Anti-stigma interventions: theory and evidence. In: Bhugra D, Bhui K, Wong SYS, Gilman SE, eds. *Oxford Textbook of Public Mental Health*. Oxford University Press; 2018: 411-420.
128. McCulloch SP, Scrivano RM. The effectiveness of mental illness stigma-reduction interventions: a systematic meta-review of meta-analyses. *Clin Psychol Rev*. 2023;100:102242. doi:[10.1016/j.cpr.2022.102242](https://doi.org/10.1016/j.cpr.2022.102242)
129. Thornicroft G, Deb T, Henderson C. Community mental health care worldwide: current status and further developments. *World Psychiatry*. 2016;15(3):276-286. doi:[10.1002/wps.20349](https://doi.org/10.1002/wps.20349)
130. Lannin DG, Bible J. Self-stigma of seeking help: a meta-analysis. In: Vogel DL, Wade NG, eds. *The Cambridge Handbook of Stigma and Mental Health*. Cambridge University Press; 2022:111-142. doi:[10.1017/9781108920995.009](https://doi.org/10.1017/9781108920995.009)
131. Lynch L, Moorhead A, Long M, Hawthorne-Steele I. The role of informal sources of help in young people's access to, engagement with, and maintenance in professional mental health care—a scoping review. *J Child Fam Stud*. 2023;32(11):3350-3365. doi:[10.1007/s10826-022-02498-5](https://doi.org/10.1007/s10826-022-02498-5)
132. Corrigan PW, Larson JE, Rüsch N. Self-stigma and the “why try” effect: impact on life goals and evidence-based practices. *World Psychiatry*. 2009;8(2):75-81. doi:[10.1002/j.2051-5545.2009.tb00218.x](https://doi.org/10.1002/j.2051-5545.2009.tb00218.x)
133. McGinty EE, Alegria M, Beidas RS, et al. The Lancet Psychiatry Commission: transforming mental health implementation research. *Lancet Psychiatry*. 2024;11(5):368-396. doi:[10.1016/S2215-0366\(24\)00040-3](https://doi.org/10.1016/S2215-0366(24)00040-3)
134. Stuart H, Sartorius N. *Paradigms Lost, Paradigms Found: Lessons Learned in the Fight Against the Stigma of Mental Illness*. 2nd ed. Oxford University Press; 2022. doi:[10.1093/med/9780197555804.001.0001](https://doi.org/10.1093/med/9780197555804.001.0001)

135. Kohn R, Ali AA, Puac-Polanco V, et al. Mental health in the Americas: an overview of the treatment gap. *Rev Panam Salud Publica*. 2018;42:e165. doi:10.26633/RPSP.2018.165
136. Shapiro E, Al-Krenawi A, Zukerman G, Melamed M, Dana Shtibi C, Korn L. Changing attitudes about mental health stigma in Israel during the COVID-19 period: a potential positive effect of the pandemic. *Soc Ment Health*. Published online September 30, 2024. doi:10.1177/21568693241276535
137. Boutron I, Tubach F, Giraudeau B, Ravaud P. Blinding was judged more difficult to achieve and maintain in nonpharmacologic than pharmacologic trials. *J Clin Epidemiol*. 2004;57(6):543-550. doi:10.1016/j.jclinepi.2003.12.010

#### SUPPLEMENT 1.

**eAppendix 1.** Protocol Amendments

**eAppendix 2.** Search Strategy

**eTable 1.** Characteristics of the Studies

**eTable 2.** Risk of Bias Within Studies

**eTable 3.** Univariable Multilevel Meta-Regression Analyses of Selected Study Characteristics for Short-Term Effects of Interventions vs Controls on Stigma-Related Attitudes

**eTable 4.** Univariable Multilevel Meta-Regression Analyses of Selected Study Characteristics for Short-Term Effects of Interventions vs Controls on Stigma-Related Knowledge

**eTable 5.** Univariable Multilevel Meta-Regression Analyses of Selected Study Characteristics for Short-Term Effects of Interventions vs Controls on General Stigma

**eTable 6.** Univariable Multilevel Meta-Regression Analyses of Selected Study Characteristics for Short-Term Effects of Interventions vs Controls Stigma-Related Behaviors

**eTable 7.** Univariable Multilevel Meta-Regression Analyses of Selected Study Characteristics for Short-Term Effects Between Active Interventions Stigma-Related Attitudes

**eTable 8.** Univariable Multilevel Meta-Regression Analyses of Selected Study Characteristics for Short-Term Effects Between Active Interventions Stigma-Related Behaviors

**eTable 9.** Univariable Multilevel Meta-Regression Analyses of Selected Study Characteristics for Short-Term Effects of Interventions vs Controls on Help-Seeking Intentions

**eTable 10.** Meta-Analytic Estimates for Secondary Outcomes (Continuous Data)

**eTable 11.** Meta-Analytic Estimates for Secondary Outcomes (Dichotomous Data)

**eFigure 1.** Forest Plot of Short-Term Effects of Interventions vs Controls on Stigma-Related Attitudes

**eFigure 2.** Forest Plot of Short-Term Effects of Interventions vs Controls on Stigma-Related Behaviors

**eFigure 3.** Forest Plot of Short-Term Effects of Interventions vs Controls on Stigma-Related Knowledge

**eFigure 4.** Forest Plot of Short-Term Effects of Interventions vs Controls on General Stigma

**eFigure 5.** Forest Plot of Short-Term Effects Between Active Interventions on Stigma-Related Attitudes

**eFigure 6.** Forest Plot of Short-Term Effects Between Active Interventions on Stigma-Related Behaviors

**eFigure 7.** Forest Plot of Short-Term Effects of Interventions vs Controls on Help-Seeking Intentions

**eFigure 8.** Funnel Plot of Short-Term Effects of Interventions vs Controls on Stigma-Related Attitudes

**eFigure 9.** Funnel Plot of Short-Term Effects of Interventions vs Controls on Stigma-Related Behaviors

**eFigure 10.** Funnel Plot of Short-Term Effects of Interventions vs Controls on Stigma-Related Knowledge

**eFigure 11.** Funnel Plot of Short-Term Effects of Interventions vs Controls on General Stigma

**eFigure 12.** Funnel Plot of Short-Term Effects Between Active Interventions on Stigma-Related Attitudes

**eFigure 13.** Funnel Plot of Short-Term Effects Between Active Interventions on Stigma-Related Behaviors

**eFigure 14.** Funnel Plot of Short-Term Effects of Interventions vs Controls on Health-Seeking Intentions

#### SUPPLEMENT 2.

**Data Sharing Statement**